
(19)

WORLD INTELLECTUAL PROPERTY ORGANIZATION

(11)Publication
number:

03057884 KR

(43)Date of publication of application:
17.07.2003

(21)Application
number: KR20030000015

(22)Date of filing: 06.01.2003

(71)Applicant:

LEADBIO INC.
ANGIOLAB INC
KIM
Jeong-Yoon
KIM
Min-Young
PARK
Eun-Kyu
CHANG
Jae-Young
KANG
Hyun-Ah

(72)Inventor:

KIM
Jeong-Yoon
KIM
Min-Young
PARK
Eun-Kyu
CHANG
Jae-Young
KANG
Hyun-Ah

(51)Int. Cl

C12N 15/14

(54) HUMAN SERUM ALBUMIN-TIMP2 FUSION PROTEIN, A POLYNUCLEOTIDE ENCODING THE SAME AND A METHOD OF PRODUCING THE HUMAN SERUM ALBUMIN-TIMP2 FUSION PROTEIN

(57) Abstract:

The present invention provides a human serum albumin-TIMP2 fusion protein having the amino acid sequence set forth in SEQ ID NO. 10, a polynucleotide encoding the same and a vector comprising the polynucleotide, a host cell transformed with the vector, a method for producing the human serum albumin-TIMP2 fusion protein and a pharmaceutical composition comprising the human serum albumin-TIMP2 fusion protein. The human serum albumin-TIMP2 fusion protein is stable and retains the activity of TIMP2, thus it can be used as a pharmaceutical composition to treat diseases related to angiogenesis and/or metastasis of cancer cells.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 July 2003 (17.07.2003)

PCT

(10) International Publication Number
WO 03/057884 A1

(51) International Patent Classification⁷: **C12N 15/14**

(21) International Application Number: **PCT/KR03/00015**

(22) International Filing Date: 6 January 2003 (06.01.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
10-2002-0001057 8 January 2002 (08.01.2002) KR

(71) Applicants (*for all designated States except US*): **LEAD-BIO INC.** [KR/KR]; Rm. 1721 Yuseong Hanjin Resort Officetel 535-5, Bongmyung-dong, Yuseong-gu, 305-301 Daejeon-city (KR). **ANGIOLAB INC** [KR/KR]; Paichai University, 439-6, Doma2-dong, Seo-gu, 302-735 Daejeon-city (KR).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **KIM, Jeong-Yoon** [KR/KR]; Chungnam National University, 220, Gung-dong, Yuseong-gu, 305-764 Daejeon-city (KR). **KIM, Min-Young** [KR/KR]; 111-801 Samsung

Pureun Apt., Jeonmin-dong, Yuseong-gu, 305-390 Daejeon-city (KR). **PARK, Eun-Kyu** [KR/KR]; 45-7, Domal-dong, Seo-gu, 302-819 Daejeon-city (KR). **CHANG, Jae-Young** [KR/KR]; Chungnam National University, 220, Gung-dong, Yuseong-gu, 305-764 Daejeon-city (KR). **KANG, Hyun-Ah** [KR/KR]; 102-1202 Lucky Hana Apt., Shinsung-dong, Yuseong-gu, 305-721 Daejeon-city (KR).


(74) Agent: **LEE, Young-Pil**; The Cheonghwa Building, 1571-18, Seocho-dong Seocho-gu, 137-874 Seoul (KR).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,

[Continued on next page]

(54) Title: **HUMAN SERUM ALBUMIN-TIMP2 FUSION PROTEIN, A POLYNUCLEOTIDE ENCODING THE SAME AND A METHOD OF PRODUCING THE HUMAN SERUM ALBUMIN-TIMP2 FUSION PROTEIN**



(57) Abstract: The present invention provides a human serum albumin-TIMP2 fusion protein having the amino acid sequence set forth in SEQ ID NO. 10, a polynucleotide encoding the same and a vector comprising the polynucleotide, a host cell transformed with the vector, a method for producing the human serum albumin-TIMP2 fusion protein and a pharmaceutical composition comprising the human serum albumin-TIMP2 fusion protein. The human serum albumin-TIMP2 fusion protein is stable and retains the activity of TIMP2, thus it can be used as a pharmaceutical composition to treat diseases related to angiogenesis and/or metastasis of cancer cells.

WO 03/057884 A1



ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI,
SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN,
GQ, GW, ML, MR, NE, SN, TD, TG).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

— *with international search report*

HUMAN SERUM ALBUMIN-TIMP2 FUSION PROTEIN, A POLYNUCLEOTIDE
ENCODING THE SAME AND A METHOD OF PRODUCING THE HUMAN
SERUM ALBUMIN-TIMP2 FUSION PROTEIN

5 Technical Field

 The present invention relates to a fusion protein, and particularly, to a human serum albumin-TIMP2 fusion protein, a polynucleotide encoding the same, a vector comprising the polynucleotide, a transformed host cell comprising the vector, a pharmaceutical composition comprising the human
10 serum albumin-TIMP2 fusion protein and a method for producing the human serum albumin-TIMP2 fusion protein.

Background Art

 Angiogenesis is the process of generating new capillary blood vessels.
15 During angiogenesis, neovascularization is tightly regulated and activation thereof occurs in embryogenic development, tissue remodeling, wound healing, and periodic cycles of corpus luteum development (Folkman and Cotran, *Int. Rev. Exp. Pathol.*, **16**, 207-248, 1976).

 During the process of angiogenesis, capillary blood vessel endothelial
20 cells start to proliferate from an existing vasculature. The endothelial cells grow very slowly as compared with other types of cells in a body. The proliferation of these cells is induced by pro-angiogenic cytokines, inflammation mediators, and activated proteolytic enzymes.

 Failure to regulate angiogenesis leads to the development of several
25 clinical syndromes or conditions. Pathological angiogenesis is involved in various diseases such as cancer in metastatic phase, arthritis, psoriasis, and retinopathy.

Not only reorganization of the blood vessel by migration, proliferation and differentiation of endothelial cells, but also degradation of an extracellular matrix is required for angiogenesis. One of the major events for inducing angiogenesis is a breakdown of the extracellular matrix before the formation of the capillary blood vessels. One of the most important enzymes which are involved in the matrix degradation is matrix metalloproteinase (MMP), a family of over 20 proteins. MMPs are endopeptidases, which degrade or proteolyze various components of the extracellular matrix such as collagen, proteoglycan, and gelatin.

10 An MMP's activity is modulated by an endogenous substance called Tissue Inhibitors of Metalloproteinases (TIMP) (Liotta and Stetler-Stevenson, *Semin. Cancer. Biol.* 1(2), 99-106, 1990; Liotta *et al.*, *Cell*, 64(2), 327-336, 1991). The proteins in the TIMP family are classified as tumor suppressor proteins and four proteins have been identified as members of this family.

15 TIMP2, one of the four identified proteins in TIMP family, is able to bind to pro- and active form of MMP-2. Since TIMP2 inhibits all the activated forms of MMPs, TIMP2 acts as a key inhibitor molecule in angiogenesis and cancer metastasis. For example, tumor cell growth and metastasis were inhibited by a gene therapy with TIMP2 in experimental animals (Hajitou *et al.*, *Cancer Res.*, 20 61, 3450-3457, 2001; Li *et al.*, *Human Gene Ther.* 12, 515-526, 2001; Sacco *et al.*, *Gene Ther.*, 8, 67-70, 2001). However, studies with TIMP2 were very limited due to a very limited amount of the protein existing in a biological system. Therefore, it is indispensable to develop a recombinant technique for overexpressing the TIMP-2 protein *in vitro*.

Although *E. coli* is a preferred host in recombinant DNA technology for producing large quantities of heterologous proteins economically, certain foreign proteins expressed in large quantities from *E. coli* are precipitated as inclusion bodies. Recovery of a biologically active protein from these
5 inclusion bodies has presented critical problems and the recovered proteins are often biologically inactive because they are folded into a three-dimensional conformation different from that of native protein. Since TIMP2 has 6-disulfide linkages, it is very complicated to refold denatured TIMP2 into its correct, biologically active conformation.

10 As a eukaryote, yeast is a suitable host organism for a high-level production of secreted soluble cytosolic proteins of human origin. Indeed, many kinds of pharmaceutically important proteins have been expressed in yeast. Yeast is able to splice out introns and transport proteins through secretory pathways as higher eukaryotes do. Especially, *Saccharomyces*
15 *cerevisiae*, the molecular and cellular biology of which has been intensively studied, has been exploited as a host for heterologous protein production since essential elements for gene expression such as strong and regulable promoters, vectors, and genetic markers are well developed (Romanos *et al.*, *Yeast*, 8, 423-488, 1992). Moreover, its use in food fermentation for
20 thousands of years proved that *S. cerevisiae* causes no harm to human beings and the processes for the production of therapeutic proteins using yeast acquired GRAS (generally recognized as safe) status. Altogether, these features make *S. cerevisiae* one of the most suitable organisms for heterologous gene expression.

Despite many advantages of yeast expression systems, a number of proteins are neither expressed in a large quantity nor secreted efficiently in yeast for unknown reasons. When human TIMP2 is expressed in yeast, for example in *S. cerevisiae*, the expression level is extremely low.

5 Human serum albumin (HSA) consisting of 585 amino acids is the most abundant protein in plasma, representing about 60 % of total plasma proteins. A major function of serum albumin is to maintain a natural osmotic pressure of plasma and to transport sparingly soluble substances throughout the body. Serum albumin also functions as a carrier of endogenous and exogenous
10 molecules, and for many years it has been thought to be devoid of any enzymatic function. However, recently, it has been found that it acts as dihydrotestosterone enolase and phospholipid cysteine peroxidase (Drmanovic *et al.*, *Anticancer Res.* 19(5B), 4113-4124, 1999; Hurst *et al.*, *Biochem J.*, 338(Pt3), 723-728, 1999). Despite these findings, exogenously administered
15 modified serum albumins, for example recombinant therapeutic proteins fused to serum albumin, are not likely to contribute significantly to the total albumin pool because of the relative abundance of albumin in plasma. Furthermore, human serum albumin is a very stable protein displaying an *in vivo* half-life of 19 days in the adult human (Sterling, K., *J. Clin. Invest.*, 30, 1228, 1957).

20

Disclosure of the Invention

It is an object of the present invention to provide a human serum albumin-TIMP2 fusion protein.

It is another object of the present invention to provide a polynucleotide
25 encoding the human serum albumin-TIMP2 fusion protein.

It is another object of the present invention to provide a vector comprising a polynucleotide encoding a human serum albumin-TIMP2 fusion protein.

It is another object of the present invention to provide a transformed host
30 cell with a vector comprising a polynucleotide encoding the human serum

albumin-TIMP2 fusion protein.

It is another object of the present invention to provide a method for producing a human serum albumin-TIMP2 fusion protein.

It is yet another object of the present invention to provide a
5 pharmaceutical composition comprising a human serum albumin-TIMP2 fusion protein.

The present invention provides a human serum albumin-TIMP2 fusion protein having the amino acid sequence of SEQ ID NO. 10. The fusion
10 protein has an activity of inhibiting the MMP enzyme activity and angiogenesis.

The molecular weight of the fusion protein is about 87.6 kDa. The fusion protein is made by fusing the carboxyl terminus of a human serum albumin to the amino terminus of TIMP2. When the fusion protein is linked to secretory signal sequence, it can be secreted to a medium more efficiently than the
15 TIMP2. Moreover, the fusion protein is more stable than TIMP2 by being fused to a human serum albumin.

The present invention also provides a polynucleotide encoding a human serum albumin-TIMP2 fusion protein having the amino acid sequence of SEQ ID NO. 10. Preferably, the polynucleotide is a polynucleotide having
20 nucleotide sequence set forth in SEQ ID NO. 3.

The present invention also provides a vector comprising a polynucleotide encoding a human serum albumin-TIMP2 fusion protein having the amino acid sequence of SEQ ID NO. 10. Preferably, the polynucleotide is
25 a polynucleotide having nucleotide sequence set forth in SEQ ID NO. 3. The vector may include any element to establish a conventional function as a vector, for example, promoter, terminator, selection marker, and origin of replication. The promoter can be constitutive or regulative, and is selected from, for example, promoters of genes for galactokinase (GAL1), uridylyltransferase
30 (GAL7), epimerase (GAL10), phosphoglycerate kinase (PGK), glyceraldehyde-3-phosphate dehydrogenase (GPD), alcohol dehydrogenase

(ADH), and the like. The vector may further include a polynucleotide encoding a secretory signal sequence to secrete the fusion protein into a medium. The polynucleotide includes a polynucleotide encoding a secretory signal sequence which is, for example, a polynucleotide encoding a human serum albumin presequence having the nucleotide sequence set forth in SEQ ID. 2. Generally, the secretory signal sequence guides heterologous proteins through the secretory pathways of a host cell, for example, a yeast cell and finally to the culture medium. Preferably, the vector is a pHSATIMP. Table 1 describes the components of pHSATIMP. FIG. 1 is a schematic representation of the pHSATIMP plasmid.

[Table 1]

	pHSATIMP
Promoter	<i>GAL10</i> promoter (SEQ ID NO. 1)
Signal sequence	HSA signal sequence (SEQ ID NO. 2)
Structural gene	HSA-TIMP2 fusion gene (SEQ ID NO. 3)
Terminator	<i>GAL7</i> terminator (SEQ ID NO. 4)
Selection marker	<i>URA3</i> gene
Origin of replication	$2 \mu m$

The present invention also provides a host cell transformed with a vector comprising a polynucleotide encoding a human serum albumin-TIMP2 fusion protein having the amino acid sequence of SEQ ID NO. 10. Preferably, the polynucleotide is a polynucleotide having nucleotide sequence set forth in SEQ ID NO. 3. The host cell for the expression of the said polynucleotide, SEQ ID NO. 3, can be any cell, including yeasts, that can be used for the heterologous gene expression. As regards yeasts, preferred genera are *Saccharomyces*, *Pichia*, *Hansenula*, *Yarrowia*, *Kluyveromyces*, and *Schizosaccharomyces*. One example of the transformed host cell is *S. cerevisiae* JY28 [a strain (*MAT pep4::HIS3 prb-1.6R can1 his3-20 ura3-52*) carrying pHSATIMP] (KCTC

10131BP).

The present invention also provides a method for producing a human serum albumin-TIMP2 fusion protein by cultivating a host cell transformed with
5 a vector comprising a polynucleotide encoding a human serum albumin-TIMP2 fusion protein having the amino acid sequence of SEQ ID NO. 10 in a suitable medium to produce the fusion protein and recovering the fusion protein. Preferably, the polynucleotide is a polynucleotide having nucleotide sequence set forth in SEQ ID NO. 3. One example of the transformed host cell is *S.*
10 *cerevisiae* JY28 [a strain (*MAT pep4::HIS3 prb-1.6R can1 his3-20 ura3-52*) carrying pHSATIMP] (KCTC 10131BP). The medium varies depending on a selected host cell, and includes a conventional medium used for cultivating the selected host cell. For example, if the host cell is *S. cerevisiae* JY28 [a strain (*MAT pep4::HIS3 prb-1.6R can1 his3-20 ura3-52*) carrying pHSATIMP] (KCTC
15 10131BP), a minimal medium containing 6.7 grams of yeast nitrogen base (without amino acids) (YNB) (Difco), 20 grams of glucose, and 20 grams of agar per liter can be used for the maintenance. The transformed host cell can be induced to produce the human HSA-TIMP2 by cultivating for 2 days at 30 °C in an induction medium composed of 10 grams of yeast extract (Difco), 20
20 grams of Bacto-peptone (Difco), 10 grams of glucose, and 20 grams of galactose per liter. The host cell transformed with the vector containing the polynucleotide, SEQ ID NO. 3, expresses and secretes the recombinant fusion protein of about 87.6 kDa in size. The recombinant fusion protein defines human TIMP2 protein fused with human serum albumin, retaining the biological
25 activity of human TIMP2 protein.

Conventional separation and purification methods for protein can be used to purify the human serum albumin-TIMP2 fusion protein of the present invention. For example, a salting out, a dialysis, an ion chromatography and an affinity chromatography can be used. When *S. cerevisiae* JY28 [a strain
30 (*MAT pep4::HIS3 prb-1.6R can1 his3-20 ura3-52*) carrying pHSATIMP] (KCTC 10131BP) was used, the final concentration of the fusion protein in culture

supernatant was about 30-50 mg per liter in a flask culture. Yeast strains including *S. cerevisiae* are known to secrete only a little amount of TIMP2, but the secretion efficiency was increased more than 50- to 100-folds by fusing TIMP2 with human serum albumin protein in the present invention.

5

The present invention also provides a pharmaceutical composition comprising a pharmaceutically effective amount of the human serum albumin-TIMP2 fusion protein and a pharmaceutically acceptable diluent or carrier. The diluent or carrier can be any material conventionally used for a pharmaceutical composition comprising a protein. The recombinant fusion protein in the present invention not only inhibits the activity of MMPs (Figure 4) but also suppresses the tube formation of human vein umbilical cells (Figure 5). Therefore, the human serum albumin-TIMP2 fusion protein of the present invention retains biological activity of the TIMP2 protein, and the fusion protein is expected to be pharmaceutically useful without any undesirable side effects because serum albumin used as a fusion partner is known to be the most abundant protein in plasma. Thus, the fusion protein can be used as an anti-angiogenic protein. In particular, the fusion protein is a potent therapeutic agent to treat diseases related to angiogenesis and/or metastasis of cancer cells and may be more useful than TIMP2 itself because of its prolonged *in vivo* stability endowed by its fusion partner, human serum albumin.

15
20

Brief Description of the Drawings

FIG. 1 is a schematic representation of the pHSATIMP plasmid.

25

FIG. 2 shows the SDS-PAGE and Western blotting results of the culture supernatants of *S. cerevisiae* JY28 (Y2805/pHSATIMP).

FIG. 3 shows the SDS-PAGE for a purified recombinant HSA-TIMP2 fusion protein.

FIG. 4 shows the inhibitory activity of the purified recombinant

HSA-TIMP2 protein on MMP-2.

FIG. 5 shows the effect of the recombinant HSA-TIMP2 on the tube formation of human umbilical vein endothelial cells (HUVECs).

5

Best mode for carrying out the invention

The following examples are intended to further illustrate the present invention. However, these examples are presented only for a better
10 understanding of the present invention without limiting its scope.

Example 1: Construction of the recombinant expression vector

In the present example, pHSATIMP containing *GAL10* promoter, HSA
15 signal sequence, HSA structural gene, TIMP2 structural gene, and *GAL7*
terminator was prepared. The gene for human serum albumin was amplified
by PCR. The template was the HSA gene in the plasmid pHSA (LeadBio, Inc)
and the primers used were SEQ ID NO. 5 (forward primer with the recognition
sequence for *EcoRI*) and SEQ ID NO. 6 (backward primer containing 15mers
20 that are complementary to the primer of SEQ ID NO. 7). The gene for human
TIMP2 was also amplified by PCR. The template was the TIMP2 gene in the
plasmid pMY2 (AngioLab, Inc) and the primers used were SEQ ID NO. 7
(forward primer) and SEQ ID NO. 8 (backward primer with the recognition
sequence for *HindIII*). The 15mers of 3' terminal sequence of the amplified

human serum albumin gene and the 15mers of 5' terminal sequence of the TIMP2 gene are complementary to each other. Thus, in-frame fusion of the human serum albumin and TIMP2 genes could be made by PCR using SEQ ID NO. 5 and SEQ ID NO. 8 as primers. The PCR product treated with restriction enzymes *EcoRI* and *HindIII* were ligated with the vector pHSA cut with *EcoRI* and *HindIII*, resulting in the recombinant vector, pHSATIMP.

Example 2: Construction of transformant

10

The plasmid, pHSATIMP prepared in Example 1 was introduced into *S. cerevisiae* Y2805 (*MAT pep4::HIS3 prb-1.6R can1 his3-20 ura3-52*) by lithium acetate method (Ito et al., *J. Bacteriol.* 153, 163-168, 1983). The selected transformants were further tested by further growing them on a synthetic complete medium without uracil. The finally selected transformant was named *S. cerevisiae* JY28 (Y2805/pHSATIMP) and deposited in KCTC (Korean Collection for Type Cultures) 10131BP on December 3, 2001.

15

Example 3: Expression of the human serum albumin and TIMP2 fusion protein

20

The transformant obtained in Example 2 was grown for 2 days at 30°C in 50 ml of YPDG medium (1% Yeast extract, 2% Proteose-peptone, 1% glucose, 2% galactose). When glucose in the medium was depleted, *GAL10* promoter was turned on by galactose and the HSA-TIMP2 fusion protein was

expressed and secreted. Culture supernatants (20 μ l) taken after 24 and 48 hours were analyzed on an SDS-PAGE gel by staining the gel with coomassie blue or after immunoblotting using a rabbit polyclonal serum directed against HSA. Figure 2 displays the SDS-PAGE and Western blotting results of the culture supernatants of *S. cerevisiae* JY28 (Y2805/pHSATIMP). Fig. 2A is a result of SDS-PAGE of the culture supernatants with Coomassie blue staining. Fig. 2B is a result of Western blotting with a rabbit polyclonal serum directed to a human serum albumin. Lanes 1, 2, 3, and 4 indicate a molecular weight marker, a supernatant(10 μ l) of a control strain (Y2805) culture after 48 hours of growth; a supernatant(10 μ l) of a JY28 (Y2805/pHSATIMP) culture after 24 hours of growth, and a supernatant(10 μ l) of a JY28 (Y2805/pHSATIMP) culture after 48 hours of growth, respectively. The HSA-TIMP2 fusion protein having the size of 87.6 kDa is clearly shown in lanes 3 and 4.

15

Example 4: Purification of the recombinant HSA-TIMP2 protein

In order to purify the HSA-TIMP2, yeast culture media of Example 3 was recovered after centrifugation at 10,000 x g for 10 min. Proteins in the supernatant were precipitated with 70% of ammonium sulfate solution. Pellets were collected by centrifugation for 30 min at 15,000 x g, and redissolved in 50 mM HEPES buffer (pH 8.0). After removal of ammonium sulfate by dialysis, the concentrated protein solution was subjected to DEAE-sepharose

(Pharmacia) column chromatography. The column was washed with 50 mM HEPES buffer, pH 8.0, and bound proteins were eluted with linear gradient of 0.1-0.5 M NaCl solution. HSA-TIMP2 was eluted at 0.24 M of NaCl, the protein was analyzed on SDS-PAGE. As shown in Figure 3, the molecular
5 size of the purified recombinant HSA-TIMP2 fusion protein was 87.6 kDa. About 20 mg of the HSA-TIMP2 was obtained from 1L of culture media.

Example 5: Effect of the recombinant HSA-TIMP2 on Matrix

10 Metalloproteinase activity

(1) Preparation of MMP

MMP-2 cDNA (GENEBANK No. XM_048244) was cloned and prepared from insect cells (Sf21 insect cell) by using a Baculovirus system.

15 The obtained MMP-2 cDNA was cloned to a pBlueBac4.5 transfer vector (Invitrogen, Cat no. V1995-20), and then transfected to Sf21 cells with a Bac-N-Blue Transfection Kit (Invitrogen, Cat no. K855-01). Sf21 cells were cultured in TNM-FH media (Sigma, St. Louis, MO, U.S.A) containing 10% fetal bovine serum at 27 °C, then harvested and re-suspended at a concentration of
20 10^7 cell/ml. The cell suspension was incubated with a virus containing the cloned gene for 1 hr at room temperature. Infected Sf21 cells were grown for 72 hrs and the medium was recovered, and the MMP-2 was purified using a gelatin-sepharose affinity column (Sigma, G5384) chromatography.

(2) Inhibition of MMP activity

In order to investigate MMP inhibition by the recombinant HSA-TIMP2 fusion protein, MMP activity was assayed by a spectrofluorometric method using Perkin-Elmer LS50B.

5 Purified MMP-2 was used after activation with 1 mM APMA before assay. The substrate for MMP-2 was MCA-Pro-Leu-Gly-Leu-Dap(Dnp)-Ala-Arg-NH₂ (Bachem, Cat. No. M-1895).

For control, 2 ml of reaction buffer (50 mM Tricine, pH 7.5, 10 mM CaCl₂, 200 mM NaCl) comprising DMSO, 10 nM of MMP-2 and 10 µM of
10 substrate was prepared in a cuvette, and fluorescence intensity was measured for 5-10 min. at room temperature with a spectrofluorometer under an excitation wavelength of 328 nm and an emission wavelength of 393 nm.

HSA-TIMP2 was added to a reaction buffer containing a substrate and MMP-2, and fluorescence intensity was measured in the same manner.

15 Figure 4 shows the inhibitory activity of the purified recombinant HSA-TIMP2 fusion protein on MMP-2. As shown in Figure 4, 7.5 µg/ml and 15 µg/ml of HSA-TIMP2 inhibited about 37% and 55% of MMP-2 activity, respectively.

20 **Example 6: Effect of the recombinant HSA-TIMP2 fusion protein on tube formation of HUVEC**

The effect of the recombinant HSA-TIMP2 fusion protein on human

endothelial cells was investigated to evaluate the biological effect of the HSA-TIMP2 of the present invention. Since MMPs are responsible for the degradation of extracellular matrix, TIMP2 is able to inhibit the formation of tubular network of vessel, which represents migration and differentiation of endothelial cell.

Blood vessel endothelial cells, human umbilical vein endothelial cells (HUVECs), were isolated from freshly obtained cords after a cesarean section according to Grants' method (Grants *et al.*, *Cell*, 58, 933-943, 1989). They were identified by immunocytochemical staining with anti-Factor VIII antibody.

HUVECs cultured on Matrigel (BD Bioscience, Bedford, MA, USA) were treated with 6.5 $\mu\text{g/ml}$ of HSA-TIMP2, and further incubated at 37 °C for 8-16 hrs. For control, the procedure was repeated with a solution without the recombinant HAS-TIMP2 fusion protein.

Fig. 5 shows the effect of the recombinant HAS-TIMP2 on a tube formation of human umbilical vein endothelial cells (HUVECs). Fig. 5A shows that a tubular network was formed as a process of neovascularization, when the HUVECs were grown on Matrigel. However, the microvascular network was disconnected when the HUVECs grown on Matrigel were treated with 6.5 $\mu\text{g/ml}$ of HSA-TIMP2 (Fig. 5B). These data show that HSA-TIMP2 is able to inhibit angiogenesis by inhibiting MMP activity.

When the area of the tubular network of HUVECs was determined using an image analysis program, Image-Pro Plus® (Media Cybernetics, USA), the tube area after treatment of HSA-TIMP2 was about 19% as compared with the untreated control. That is, the tube formation was inhibited by 81% with 6.5

μg/ml of HSA-TIMP2 of the present invention.

Industrial Applicability

5 According to the human serum albumin-TIMP2 fusion protein of the present invention, it retains the biological activity of the TIMP2 and can be used as a pharmaceutically active component without any undesirable side effects.

10 According to the polynucleotide and the vector comprising the same of the present invention, it is able to express the human serum albumin-TIMP2 fusion protein of the invention.


According to the pharmaceutical composition of the present invention, it can be used as an anti-angiogenic protein or as a potent therapeutic agent to treat diseases related to angiogenesis and/or metastasis of cancer cells.

15 According to the method of the present invention, the human serum albumin-TIMP2 fusion protein of the present invention can be produced on a large scale by using a transformed host cell.

Applicant's or agent's file reference	International application No.
--	-------------------------------

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page <u>6</u> , line <u>22</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution <p align="center">Korean Collection for Type Cultures</p>	
Address of depositary institution (including postal code and country) <p align="center">Korea Research Institute of Bioscience and Biotechnology (KRIBB) #52, Oun-dong, Yusong-ku, Taejon 305-333 Rep. of Korea</p>	
Date of deposit <p align="center">03 December 2001</p>	Accession Number <p align="center">KCTC 10131BP</p>
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	
<p align="center">For receiving Office use only</p> <div><input type="checkbox"/> This sheet was received with the international application</div> <div>Authorized officer </div>	<p align="center">For International Bureau use only</p> <div><input type="checkbox"/> This sheet was received by the International Bureau on:</div> <div>Authorized officer</div>

Form PCT/RO/134 (July 1998)

What is claimed is:

1. A polypeptide having the amino acid sequence set forth in SEQ ID NO. 10.
- 5 2. A polynucleotide encoding the polypeptide according to claim 1.
3. The polynucleotide of claim 2, wherein the polynucleotide
10 is a polynucleotide having the nucleotide sequence set forth in SEQ ID NO. 3.
4. A vector comprising a polynucleotide encoding the polypeptide having the amino acid sequence set forth in SEQ ID NO. 10.
- 15 5. The vector of claim 4, wherein the polynucleotide is a polynucleotide having the nucleotide sequence set forth in SEQ ID NO. 3.
- 20 6. The vector of claim 4, further comprising a polynucleotide encoding a secretory signal sequence for extracellular secretion of a protein.
7. The vector of claim 6, wherein the secretory signal
25 sequence comprises a polynucleotide encoding a human serum albumin presequence
8. The vector of claim 4, wherein the vector is pHSATIMP.
- 30 9. A host cell transformed with a vector according to any one of claims 4 through 8.
10. The host cell of claim 9, wherein the host cell is a yeast.

11. The host cell of claim 9, wherein the host cell is *S. cerevisiae* JY28 (KCTC 10131BP).


5 12. A method for producing a human serum albumin-TIMP2 fusion protein comprising cultivating the transformed host cell according to claim 9 in a suitable medium to produce the human serum albumin-TIMP2 fusion protein and recovering the human serum albumin-TIMP2 fusion protein.

10

13. A pharmaceutical composition comprising a pharmaceutically effective amount of a human serum albumin-TIMP2 fusion protein having the amino acid sequence set forth in SEQ ID. 10 and a pharmaceutically acceptable diluent or carrier.


1/6

Fig. 1




2/6

Fig. 2




3/6

Fig. 3




4/6

Fig. 4




5/6

Fig. 5A



6/6

Fig. 5B



<110> ANGIOLAB, INC.
LeadBio, Inc.

<120> HUMAN SERUM ALBUMIN-TIMP2 FUSION PROTEIN, A POLYNUCLEOTIDE
5 SERUM ENCODING THE SAME AND A METHOD OF PRODUCING THE HUMAN
ALBUMIN-TIMP2 FUSION PROTEIN

<160> 10

10 <170> KopatentIn 1.71

<210> 1
<211> 507
15 <212> DNA
<213> GAL10 promoter

<400> 1
atcgcttgcg tgattaatta cccagaaat aaggctaaaa aactaatcg attatcatcc 60
20 tatgggtgtt aattgattc gttcattga aggtttgtgg ggccagggtt ctgccaatt 120
ttctcttca taaccataaa agctagtatt gtagaatctt tattgttcgg accagtgcgg 180
25 cgcgaggcac atctgogttt caggaacgcg accggtgaag acgaggacgc acggaggaga
240
gtcttccttc ggagggtgt caccgcctcg gcggcttcta atccgtactt caataatgca 300
30 atgagcagtt aagcgtatta ctgaaagttc caaagagaag gtttttttag gctaagataa 360
tggggctctt tacatttcca caacatataa gtaagattag atatggatat gtatatggat 420
atgtatatgg tgtaatgcc atgtaatatg attattaac ttcttgcgt ccatcaaaa 480
35 aaaaagtaag aattttgaa aattcaa 507

<210> 2
40 <211> 72
<212> DNA
<213> HSA pre signal

<400> 2
45 atgaagtggg taaccttat ttcccttctt ttctcttta gctoggctta ttccaggggt 60
gtgtttogtc ga 72

50 <210> 3
<211> 2340
<212> DNA
<213> Artificial Sequence

<220>

<223> HSA-TIMP2 fusion gene

5 <400> 3
 gatgcacaca agagttaggt tgctcatcgg tttaaagatt tgggagaaga aaatttcaaa 60
 gcttggtgt tgattgcctt tgctcagtat ctccagcagt gtccatttga agatcatgta 120
 10 aaattagtca atgaagtaac tgaatttga aaaacatgtg ttgctgatga gtcagctgaa 180
 aattgtgaca aatcacttca taccctttt ggagacaaat tatgcacagt tgcaactctt 240
 cgtgaaacct atggtgaaat ggctgactgc tgtgcaaaac aagaacctga gagaaatgaa 300
 15 tgcttcttgc aacacaaaga tgacaacca aacctcccc gattggtgag accagaggtt 360
 gatgtgatgt gcaactgctt tcatgacaat gaagagacat tttgaaaaa atacttata 420
 20 gaaattgcc aagacatcc ttactttat gcccgggaac tccttttctt tgctaaaagg 480
 tataaagctg cttttacaga atgttgccaa gctgctgata aagctgcctg cctgttgcca 540
 aagctcgatg aactcggga tgaagggaag gcttcgtctg ccaaacagag actcaagtgt 600
 25 gccagtctcc aaaaatttgg agaaagagct ttcaaagcat gggcggtggc tcgcctgagc 660
 cagagatttc ccaaagctga gtttcagaa gtttccaagt tagtgacaga tcttaccaaa 720
 30 gtccacacgg aatgctgcca tggagatctg ctgaaatgtg ctgatgacag ggccggacct 780
 gccaaagata tctgtgaaaa tcaagattcg atctccagta aactgaagga atgctgtgaa 840
 aaacctctgt tggaaaaatc ccaactgcatt gccgaagtgg aaaatgatga gatgcctgct 900
 35 gacttgcctt cattagctgc tgattttgtt gaaagtaagg atgttgcaa aaactatgct 960
 gaggcaaagg atgtcttctt gggcatgttt ttgtatgaat atgcaagaag gcatcctgat 1020
 40 tactctgtcg tgctgctgct gagacttgc aagacatatg aaaccactct agagaagtgc 1080
 tgtccgctg cagatcctca tgaatgctat gccaaagtgt tgatgaatt taaacctctt 1140
 gtggaagagc ctgagaattt aatcaaacia aattgtgagc ttttgagca gcttgagag 1200
 45 taaaattcc agaatgcgt attagttcgt tacaccaaga aagtaccca agtgtcaact 1260
 ccaactcttg tagaggcttc aagaacctta gaaaagtgg gcagcaaagt ttgtaaacat 1320
 50 cctgaagcaa aaagaatgcc ctgtgcagaa gactatctat ccgtggtcct gaaccagtta 1380
 tgtgtgttc atgagaaaac gccagtaagt gacagagica ccaaatgctg cacagaatcc 1440
 ttggtgaaca ggccaccatg ctttcagct ctggaagtgc atgaaacata cgttccaaa 1500

gagtttaatg ctgaaacatt caccttccat gcagatatat gcacacttgc tgagaaggag 1560
 agacaaatca agaaacaaac tgcacttgtt gagctcgtga aacacaagcc caaggcaaca
 5 1620
 aaagagcaac tgaaagctgt tatggatgat ttgcagctt ttgtagagaa gtgctgcaag 1680
 gctgacgata aggagacctg cttgcgcgag gagggtaaaa aacttgtgc tgcaagtcaa 1740
 10 gctgcottag gcttatgcag ctgctccccg gtgcacccgc aacaggcgtt ttgcaatgca 1800
 gatgtagtga tcagggccaa agcggtcagt gagaaggaag tggactctgg aaacgacatt 1860
 15 tatggcaacc ctatcaagag gatccagtat gagatcaagc agataaagat gttcaaaggg 1920
 cctgagaagg atatagagtt tatctacacg gccccctcct cggcagtgtg tggggtctcg 1980
 ctggacgttg gaggaagaa ggaatatctc attgcaggaa aggccgaggg ggacggcaag
 20 2040
 atgcacatca ccctctgtga cttcatcgtg ccctgggaca cctgagcac caccagaag 2100
 aagagcctga accacaggta ccagatgggc tgcgagtga agatcacgcg ctgccccatg
 25 2160
 atcccgctgt acatctctc cccggacgag tgcctctgga tggactgggt cacagagaag 2220
 aacatcaacg ggcaccagcg caagttctc gcctgcatca agagaagtga cggctcctgt 2280
 30 gcggtgtacc gcggcggggc gccccccaag caggagtctc tcgacatga ggaccataa
 2340
 35 2340
 <210> 4
 <211> 528
 <212> DNA
 40 <213> GAL7 terminator
 <400> 4
 taatgtatt ctagtatgt aagagtggc cttccataa aaaaaaaaaa aaagaaaaa 60
 45 gaatttagg aatacaatgc agcttgaag taaaatctgg aatattcata tgcacacaac 120
 ttottatgct tataaaagca ctaatgcctg aatttatgtt gaaaatatgt gtcacaaata 180
 aagaaactgt gacatctgac acattccac ttattgaca agaatagaat ttctttaagt 240
 50 ttccctcta gattattat ttcaaattt taggcctctg tgaagttat tacgtagaaa 300
 ttctacgat agttattagt cctaattgga tgttcagca aggtcattg tcggtgtcgt 360

tatcgagctt ggcactggcc gtcgtttac aacgtcgtga ctgggaaaac cctggcgta 420

cccaactaa tcgcottgca gcacatcccc cottgccag ctggcgtaat agcgaagagg 480

5 cccgcaccga tcgcccttc caacagttgc gcagcctgaa tggcgaat 528

<210> 5
<211> 30
10 <212> DNA
<213> Artificial Sequence

<220>
<223> HSA cloning forward primer
15

<400> 5
cacgaattcg gcacaatgaa gtgggtaacc 30

20 <210> 6
<211> 33
<212> DNA
<213> Artificial Sequence

25 <220>
<223> HSA cloning reward primer

30 <400> 6
cggggagcag ctgcataagc ctaaggcagc ttg 33

<210> 7
35 <211> 18
<212> DNA
<213> Artificial Sequence

<220>
40 <223> TIMP2 cloning forward primer

<400> 7
45 tgcagctgct ccccggtg 18

<210> 8
<211> 20
<212> DNA
50 <213> Artificial Sequence

<220>
<223> TIMP2 cloning reward primer

<400> 8
cggtgaagct ttgcttatgg

20

5

<210> 9
<211> 6
<212> PRT
<213> Artificial Sequence

10

<220>
<223> substrate for MMP-2

15

<220>
<221> MOD_RES
<222> (1)
<223> (7-methoxycoumarin-4-yl)acetyl Pro

20

<220>
<221> MOD_RES
<222> (4)
<223> N-3-(2,4-dinitrophenyl)-L-2,3-diaminopropionyl Leu

25

<220>
<221> MOD_RES
<222> (6)
<223> aminated Arg

30

<400> 9
Pro Leu Gly Leu Ala Arg

35

1 5

<210> 10
<211> 779
<212> PRT
<213> Artificial

40

<400> 10
Asp Ala His Lys Ser Glu Val Ala His Arg Phe Lys Asp Leu Gly Glu

45

1 5 10 15
Glu Asn Phe Lys Ala Leu Val Leu Ile Ala Phe Ala Gln Tyr Leu Gln
20 25 30

50

Gln Cys Pro Phe Glu Asp His Val Lys Leu Val Asn Glu Val Thr Glu
35 40 45

Phe Ala Lys Thr Cys Val Ala Asp Glu Ser Ala Glu Asn Cys Asp Lys
50 55 60

Ser Leu His Thr Leu Phe Gly Asp Lys Leu Cys Thr Val Ala Thr Leu
 65 70 75 80

5 Arg Glu Thr Tyr Gly Glu Met Ala Asp Cys Cys Ala Lys Gln Glu Pro
 85 90 95

Glu Arg Asn Glu Cys Phe Leu Gln His Lys Asp Asp Asn Pro Asn Leu
 100 105 110

10 Pro Arg Leu Val Arg Pro Glu Val Asp Val Met Cys Thr Ala Phe His
 115 120 125

Asp Asn Glu Glu Thr Phe Leu Lys Lys Tyr Leu Tyr Glu Ile Ala Arg
 15 130 135 140

Arg His Pro Tyr Phe Tyr Ala Pro Glu Leu Leu Phe Phe Ala Lys Arg
 145 150 155 160

20 Tyr Lys Ala Ala Phe Thr Glu Cys Cys Gln Ala Ala Asp Lys Ala Ala
 165 170 175

Cys Leu Leu Pro Lys Leu Asp Glu Leu Arg Asp Glu Gly Lys Ala Ser
 180 185 190

25 Ser Ala Lys Gln Arg Leu Lys Cys Ala Ser Leu Gln Lys Phe Gly Glu
 195 200 205

Arg Ala Phe Lys Ala Trp Ala Val Ala Arg Leu Ser Gln Arg Phe Pro
 30 210 215 220

Lys Ala Glu Phe Ala Glu Val Ser Lys Leu Val Thr Asp Leu Thr Lys
 225 230 235 240

35 Val His Thr Glu Cys Cys His Gly Asp Leu Leu Glu Cys Ala Asp Asp
 245 250 255

Arg Ala Asp Leu Ala Lys Tyr Ile Cys Glu Asn Gln Asp Ser Ile Ser
 260 265 270

40 Ser Lys Leu Lys Glu Cys Cys Glu Lys Pro Leu Leu Glu Lys Ser His
 275 280 285

Cys Ile Ala Glu Val Glu Asn Asp Glu Met Pro Ala Asp Leu Pro Ser
 45 290 295 300

Leu Ala Ala Asp Phe Val Glu Ser Lys Asp Val Cys Lys Asn Tyr Ala
 305 310 315 320

50 Glu Ala Lys Asp Val Phe Leu Gly Met Phe Leu Tyr Glu Tyr Ala Arg
 325 330 335

Arg His Pro Asp Tyr Ser Val Val Leu Leu Leu Arg Leu Ala Lys Thr
 340 345 350

Tyr Glu Thr Thr Leu Glu Lys Cys Cys Ala Ala Ala Asp Pro His Glu
 355 360 365

5 Cys Tyr Ala Lys Val Phe Asp Glu Phe Lys Pro Pro Val Glu Glu Pro
 370 375 380

Gln Asn Leu Ile Lys Gln Asn Cys Glu Leu Phe Glu Gln Leu Gly Glu
 385 390 395 400

10 Tyr Lys Phe Gln Asn Ala Leu Leu Val Arg Tyr Thr Lys Lys Val Pro
 405 410 415

Gln Val Ser Thr Pro Thr Leu Val Glu Val Ser Arg Asn Leu Gly Lys
 15 420 425 430

Val Gly Ser Lys Cys Cys Lys His Pro Glu Ala Lys Arg Met Pro Cys
 435 440 445

20 Ala Glu Asp Tyr Leu Ser Val Val Leu Asn Gln Leu Cys Val Leu His
 450 455 460

Glu Lys Thr Pro Val Ser Asp Arg Val Thr Lys Cys Cys Thr Glu Ser
 25 465 470 475 480

Leu Val Asn Arg Arg Pro Cys Phe Ser Ala Leu Glu Val Asp Glu Thr
 485 490 495

Tyr Val Pro Lys Glu Phe Asn Ala Glu Thr Phe Thr Phe His Ala Asp
 30 500 505 510

Ile Cys Thr Leu Ser Glu Lys Glu Arg Gln Ile Lys Lys Gln Thr Ala
 515 520 525

35 Leu Val Glu Leu Val Lys His Lys Pro Lys Ala Thr Lys Glu Gln Leu
 530 535 540

Lys Ala Val Met Asp Asp Phe Ala Ala Phe Val Glu Lys Cys Cys Lys
 545 550 555 560

40 Ala Asp Asp Lys Glu Thr Cys Phe Ala Glu Glu Gly Lys Lys Leu Val
 565 570 575

Ala Ala Ser Gln Ala Ala Leu Gly Leu Cys Ser Cys Ser Pro Val His
 45 580 585 590

Pro Gln Gln Ala Phe Cys Asn Ala Asp Val Val Ile Arg Ala Lys Ala
 595 600 605

50 Val Ser Glu Lys Glu Val Asp Ser Gly Asn Asp Ile Tyr Gly Asn Pro
 610 615 620

Ile Lys Arg Ile Gln Tyr Glu Ile Lys Gln Ile Lys Met Phe Lys Gly
 625 630 635 640

Pro Glu Lys Asp Ile Glu Phe Ile Tyr Thr Ala Pro Ser Ser Ala Val
645 650 655

5 Cys Gly Val Ser Leu Asp Val Gly Gly Lys Lys Glu Tyr Leu Ile Ala
660 665 670

Gly Lys Ala Glu Gly Asp Gly Lys Met His Ile Thr Leu Cys Asp Phe
675 680 685

10 Ile Val Pro Trp Asp Thr Leu Ser Thr Thr Gln Lys Lys Ser Leu Asn
690 695 700

His Arg Tyr Gln Met Gly Cys Glu Cys Lys Ile Thr Arg Cys Pro Met
15 705 710 715 720

Ile Pro Cys Tyr Ile Ser Ser Pro Asp Glu Cys Leu Trp Met Asp Trp
725 730 735

20 Val Thr Glu Lys Asn Ile Asn Gly His Gln Ala Lys Phe Phe Ala Cys
740 745 750



Ile Lys Arg Ser Asp Gly Ser Cys Ala Trp Tyr Arg Gly Ala Ala Pro
755 760 765

25 Pro Lys Gln Glu Phe Leu Asp Ile Glu Asp Pro
770 775

30

INTERNATIONAL SEARCH REPORT

 International application No.
 PCT/KR03/00015

A. CLASSIFICATION OF SUBJECT MATTER IPC7 C12N 15/14 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) A01N 37/18 : C07K 14/475 : C12N 15/00, 15/11, 15/85, 15/63 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Korean Patents and Applications for Inventions since 1975 Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) PubMed, Delphion, PAJ		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Nomura N, Matsubara N, Horinouchi S, Beppu T. "Secretion by <i>Saccharomyces cerevisiae</i> of human apolipoprotein E as a fusion to serum albumin." <i>Biosci Biotechnol Biochem.</i> , vol.59(3):532-4, March 1995 See the whole document	1-13
Y	Rajan SS, Lackland H, Stein S, Denhardt DT. "Presence of an N-terminal polyhistidine tag facilitates stable expression of an otherwise unstable N-terminal domain of mouse tissue inhibitor of metalloproteinase-1 in <i>Escherichia coli</i> ." <i>Protein Expr Purif.</i> , vol.13(1):67-72, June 1998 See the abstract	1-13
A	Cockett MI, Bebbington CR, Yarranton GT. "High level expression of tissue inhibitor of metalloproteinases in Chinese hamster ovary cells using glutamine synthetase gene amplification." <i>Biotechnology (NY)</i> , vol.8(7):662-7, July 1990 See the abstract	1-10
A	US 5,595,885A (The USA Department of Health and Human Services) Jan 1997 See the abstract	13
A	US 5,643,752A (Incyte Pharmaceuticals, Inc.) July 1997 See the abstract	1-12
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input checked="" type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 26 MARCH 2003 (26.03.2003)		Date of mailing of the international search report 26 MARCH 2003 (26.03.2003)
Name and mailing address of the ISA/KR  Korean Intellectual Property Office 920 Dunsan-dong, Seo-gu, Daejeon 302-701, Republic of Korea Facsimile No. 82-42-472-7140		Authorized officer AHN, Mi Jung Telephone No. 82-42-481-5593 

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/KR03/00015

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5,595,885A	Jan 1997	None	
US 5,643,752A	July 1997	WO 9618725	June 1996